

# Induced subgraphs and subtournaments

Paul Seymour, Princeton University

joint with Maria Chudnovsky

## Theorem

- (Robertson, S., 1986) *For every planar graph  $H$  there exists  $k$  such that every graph with no  $H$ -minor has tree-width at most  $k$ .*

## Theorem

- (Robertson, S., 1986) For every planar graph  $H$  there exists  $k$  such that every graph with no  $H$ -minor has tree-width at most  $k$ .
- (Robertson, S., Thomas 1994; Diestel, Gorbunov, Jensen, Thomassen 1999) For every simple planar graph  $H$  with  $n$  vertices, taking  $k = 2^{O(n^5)}$  works.

## Theorem

- (Robertson, S., 1986) For every planar graph  $H$  there exists  $k$  such that every graph with no  $H$ -minor has tree-width at most  $k$ .
- (Robertson, S., Thomas 1994; Diestel, Gorbunov, Jensen, Thomassen 1999) For every simple planar graph  $H$  with  $n$  vertices, taking  $k = 2^{O(n^5)}$  works.
- (Leaf, S., 2012) For every simple planar graph  $H$  with  $n$  vertices, taking  $k = 2^{O(n \log(n))}$  works.

## Theorem (Robertson, S., 1986)

*For every planar graph  $H$  there exists  $k$  such that every graph with no  $H$ -minor has tree-width at most  $k$ .*

## Theorem (Robertson, S., 1986)

*For every planar graph  $H$  there exists  $k$  such that every graph with no  $H$ -minor has tree-width at most  $k$ .*

*For every non-planar graph  $H$  there is no such  $k$ .*

## Theorem (Robertson, S., 1986)

*For every planar graph  $H$  there exists  $k$  such that every graph with no  $H$ -minor has tree-width at most  $k$ .*

*For every non-planar graph  $H$  there is no such  $k$ .*

Equivalently:

## Theorem

*A minor ideal has bounded treewidth if and only if some planar graph is not in it.*

## Theorem (Robertson, S., 1986)

*For every planar graph  $H$  there exists  $k$  such that every graph with no  $H$ -minor has tree-width at most  $k$ .*

*For every non-planar graph  $H$  there is no such  $k$ .*

Equivalently:

## Theorem

*A minor ideal has bounded treewidth if and only if some planar graph is not in it.*

Equivalently:

## Theorem

*A minor ideal has bounded treewidth if and only if it does not include the ideal of all planar graphs.*



## Theorem

*A minor ideal has bounded pathwidth if and only if it does not include the ideal of all forests.*

## Theorem

*A minor ideal has bounded pathwidth if and only if it does not include the ideal of all forests.*

$I(H)$  is the ideal of all graphs each component of which is a minor of  $H$ .

## Theorem

*If  $H$  is planar, a minor ideal has bounded  $f_1$  if and only if it does not include  $I(H)$  ( $f_1$  is the minimum number of vertices whose deletion leaves a graph with no  $H$  minor)*

## Theorem

*A minor ideal has bounded pathwidth if and only if it does not include the ideal of all forests.*

$I(H)$  is the ideal of all graphs each component of which is a minor of  $H$ .

## Theorem

*If  $H$  is planar, a minor ideal has bounded  $f_1$  if and only if it does not include  $I(H)$  ( $f_1$  is the minimum number of vertices whose deletion leaves a graph with no  $H$  minor)*

## Theorem

*A minor ideal has bounded  $f_2$  if and only if it does not include  $I(K_5)$  or  $I(K_{3,3})$  ( $f_2$  is the minimum  $k$  such that deleting some  $k$  vertices leaves a graph of genus at most  $k$ ).*

## Theorem

*A minor ideal has*

- *bounded  $f_3$  iff it does not include the ideal of all graphs that are subgraphs of a path*
- *bounded  $f_4$  iff it does not include the ideal of all stars*
- *bounded  $f_5$  iff it does not include the ideal of all graphs with crossing number at most one*
- *bounded  $f_6$  iff it does not include the ideal of all graphs.*

What about induced subgraphs?

What about induced subgraphs?

## Theorem

*An induced subgraph ideal has bounded clique number iff it does not include {cliques}.*

What about induced subgraphs?

### Theorem

*An induced subgraph ideal has bounded clique number iff it does not include {cliques}.*

### Theorem (Ramsey's theorem)

*An induced subgraph ideal is finite iff it includes neither of {cliques}, {graphs with no edges}.*

What about induced subgraphs?

### Theorem

*An induced subgraph ideal has bounded clique number iff it does not include {cliques}.*

### Theorem (Ramsey's theorem)

*An induced subgraph ideal is finite iff it includes neither of {cliques}, {graphs with no edges}.*

### Theorem

*An induced subgraph ideal has bounded maximum degree iff*



What about induced subgraphs?

### Theorem

*An induced subgraph ideal has bounded clique number iff it does not include {cliques}.*

### Theorem (Ramsey's theorem)

*An induced subgraph ideal is finite iff it includes neither of {cliques}, {graphs with no edges}.*

### Theorem

*An induced subgraph ideal has bounded maximum degree iff it includes neither of {cliques}, {stars}.*

What about induced subgraphs?

### Theorem

*An induced subgraph ideal has bounded clique number iff it does not include {cliques}.*

### Theorem (Ramsey's theorem)

*An induced subgraph ideal is finite iff it includes neither of {cliques}, {graphs with no edges}.*

### Theorem

*An induced subgraph ideal has bounded maximum degree iff it includes neither of {cliques}, {stars}.*

Are there any more?

# Tournaments

Possible containment relations:

- subtournament (not wqo)
- topological containment (ie subdivision) (not wqo)
- immersion (wqo)
- butterfly minor (wqo?? – open)
- strong minor (wqo - Kim).

Tournament  $G$  has **cutwidth** at most  $c$  if  $V(G)$  can be ordered  $\{v_1, \dots, v_n\}$  such that for each  $i$ , there are at most  $c$  edges from  $\{v_{i+1}, \dots, v_n\}$  to  $\{v_1, \dots, v_i\}$ .

Tournament  $G$  has **cutwidth** at most  $c$  if  $V(G)$  can be ordered  $\{v_1, \dots, v_n\}$  such that for each  $i$ , there are at most  $c$  edges from  $\{v_{i+1}, \dots, v_n\}$  to  $\{v_1, \dots, v_i\}$ .

## Theorem

*For every tournament  $H$  there exists  $c$  such that every tournament in which  $H$  cannot be immersed has cutwidth at most  $c$ .*

Tournament  $G$  has **cutwidth** at most  $c$  if  $V(G)$  can be ordered  $\{v_1, \dots, v_n\}$  such that for each  $i$ , there are at most  $c$  edges from  $\{v_{i+1}, \dots, v_n\}$  to  $\{v_1, \dots, v_i\}$ .


## Theorem

*For every tournament  $H$  there exists  $c$  such that every tournament in which  $H$  cannot be immersed has cutwidth at most  $c$ .*

Equivalently:


## Theorem

*An immersion ideal of tournaments has bounded cutwidth iff it does not include  $\{\text{tournaments}\}$ .*



## Theorem

*For every tournament  $H$  of type  $A_0$ , there exists  $c$  such that every tournament  $G$  not containing  $H$  as a subtournament can be ordered such that the maximum backdegree is at most  $c$ .*



## Theorem


*For every tournament  $H$  of type  $A_0$ , there exists  $c$  such that every tournament  $G$  not containing  $H$  as a subtournament can be ordered such that the maximum backdegree is at most  $c$ .*


Equivalently:

## Theorem

*A subtournament ideal has bounded backdegree iff it does not include  $\{type A_0\}$ .*







## Theorem

*A subtournament ideal has bounded cutwidth iff it does not include  $\{\text{type } A_0\}$ ,  $\{\text{type } B\}$  or  $\{\text{type } B'\}$ .*

## Theorem (Fradkin, S.)

*For every tournament  $H$  there exists  $c$  such that every tournament that does not contain  $H$  topologically has pathwidth at most  $c$ .*

## Theorem (Fradkin, S.)

*For every tournament  $H$  there exists  $c$  such that every tournament that does not contain  $H$  topologically has pathwidth at most  $c$ .*

Equivalently:

## Theorem

*A topological containment ideal of tournaments has bounded pathwidth iff it does not include {tournaments}.*

## Theorem (Fradkin, S.)

*For every tournament  $H$  there exists  $c$  such that every tournament that does not contain  $H$  topologically has pathwidth at most  $c$ .*


Equivalently:

## Theorem


*A topological containment ideal of tournaments has bounded pathwidth iff it does not include {tournaments}.*

## Theorem (Kim, S.)

*A strong minor ideal of tournaments has bounded pathwidth iff it does not include {tournaments}.*



TYPE A




TYPE A

## Theorem


*A subtournament ideal has bounded pathwidth iff it does not include {type A}, {type B} or {type B'}.*

# Tournaments under subtournament containment






# Tournaments under subtournament containment




## Theorem

*A subtournament ideal has bounded length backedges iff it does not include {type C}.*

# Circular interval tournaments




# Circular interval tournaments




## Theorem

*A tournament is a circular interval tournament iff it contains neither of the tournaments above.*




## Theorem (Gaku Liu)

*A subtournament ideal consists of blowups of circular interval tournaments by tournaments with bounded length backedges iff it does not include {type D}, {type D'}.*




## Theorem

*A subtournament ideal consists of tournaments orderable such that each component of long backedges is a path iff it does not include  $\{type E\}, \{type E'\}$ .*



## Theorem

*A subtournament ideal consists of boundedly fuzzy circular interval tournaments iff it does not include {type  $G$ }, {type  $G'$ }.*



TYPE F

## Problem

A subtournament ideal consists of ??? iff it does not include {type F}.

# Induced subgraphs revisited

A finite set of graphs is **heroic** if there exists  $c$  such that every graph with no induced subgraph in the set has chromatic number at most  $c$ .

## Theorem

*Every heroic set contains*

- *a disjoint union of cliques*
- *a complete multipartite graph*
- *a forest*
- *a graph whose complement is a forest.*



# Induced subgraphs revisited

A finite set of graphs is **heroic** if there exists  $c$  such that every graph with no induced subgraph in the set has cochromatic number at most  $c$ .

## Theorem

*Every heroic set contains*

- *a disjoint union of cliques*
- *a complete multipartite graph*
- *a forest*
- *a graph whose complement is a forest.*

## Conjecture: Gyárfás' 1975; Sumner 1981

For every clique  $K$  and forest  $F$ , there is a constant  $c$  such that every graph containing neither of  $K, F$  as an induced subgraph has chromatic number at most  $c$ .

$G$  has **splitness** at most  $k$  if  $V(G)$  can be partitioned into  $X, Y$ , where  $G|X$  has no clique of size  $k + 1$  and  $G|Y$  has no stable set of size  $k + 1$ .

$G$  has **splitness** at most  $k$  if  $V(G)$  can be partitioned into  $X, Y$ , where  $G|X$  has no clique of size  $k + 1$  and  $G|Y$  has no stable set of size  $k + 1$ .

## Theorem

*An induced subgraph ideal has bounded splitness iff it does not include {disjoint unions of cliques}, {complete multipartite graphs}.*

$G$  has **splitness** at most  $k$  if  $V(G)$  can be partitioned into  $X, Y$ , where  $G|X$  has no clique of size  $k + 1$  and  $G|Y$  has no stable set of size  $k + 1$ .

## Theorem


*An induced subgraph ideal has bounded splitness iff it does not include {disjoint unions of cliques}, {complete multipartite graphs}.*

## Corollary


*If Gyárfas' conjecture is true, then every set of graphs containing one of each of the four types is heroic.*



ONE EDGE



STAR



SEMISTAR

## Theorem

*For every star  $H_1$  and star complement  $H_2$ , there exists  $c$  such that if  $G$  contains neither of  $H_1, H_2$  as an induced subgraph then one of  $G, \overline{G}$  has maximum degree at most  $c$ .*

## Theorem (Reed)

*For every one-edge graph  $H_1$  and clique  $H_2$ , there exists  $c$  such that if  $G$  contains neither of  $H_1, H_2$  as an induced subgraph then  $G$  is “almost complete multipartite with at most  $c$  parts”.*

## Theorem (Reed)

*For every one-edge graph  $H_1$  and clique  $H_2$ , there exists  $c$  such that if  $G$  contains neither of  $H_1, H_2$  as an induced subgraph then  $G$  is “almost complete multipartite with at most  $c$  parts”.*

## Theorem (Norine, Reed)

*For every semistar  $H_1$  and semistar complement  $H_2$ , there exists a structure theorem.*